BUCKEYE HEALTH PLAN

MEDICAID ANNUAL HEDIS RATES 2022-2023

| Medicaid HEDIS® Measures | 2022 (MY2021) | 2023 (MY2022) | Goal |
|--|------------------|------------------|----------------|
| Initiation & Engagement of Substance Use Disorder Treatment, Initiation Total | 50.89% | ↑ 52.40% | 45.51% |
| Initiation & Engagement of Substance Use Disorder Treatment, Engagement 18-64 years | 21.59% | ^ 28.65% | 27.42% |
| Follow-up after Hospitalization for Mental Illness-7 Day Follow-up Total | 42.85% | ↓ 34.65% | 41.67% |
| Follow-up after Hospitalization for Mental Illness-30 Day Follow-up Total | 64.44% | ↓ 55.75% | 62.06% |
| Antidepressant Med Follow up - Acute Phase | 59.25% | ↓ 57.96% | 64.90% |
| Antidepressant Med Follow up - Continuation Phase | 43.04% | ↓ 40.75% | 47.70% |
| Follow-up After ED visit for Mental Illness - 7 day | 42.85% | ↑ 51.35% | 46.08% |
| Follow-up After ED visit for Mental Illness - 30 day | 64.44% | ↓ 63.93% | 58.24% |
| Follow-up After ED Visit for Substance Abuse -30 Day follow-up | 69.87% | ↓ 42.05% | 60.71% |
| Follow-up After ED Visit for Substance Abuse -7 Day follow-up | 51.10% | ↓ 28.95% | 40.03% |
| Use of Opioids at High Dosage (lower is better) | 1.37% | ↓ 1.15% | 2.04% |
| Use of Opioids from Multiple Pharmacies & Multiple Prescribers - Multiple Pharmacies (lower is better) | 3.02% | ↓ 2.46% | 1.27% |
| Use of Opioids from Multiple Pharmacies & Multiple Prescribers - Multiple Prescribers (lower is better) | 23.99% | ↓ 23.55% | 15.24% |
| Use of Opioids from Multiple Pharmacies & Multiple Prescribers - Multiple Pharmacies & | 2.08% | ↓ 1.85% | 0.74% |
| Prescribers (lower is better) Risk of Continued Opioid Use >= 15 days Total (lower is better) | 4.400/ | ↑ 6 00% | 2 610/ |
| Risk of Continued Opioid Use >= 15 days Total (lower is better) Risk of Continued Opioid Use >= 31 days Total (lower is better) | 4.40% 2.17% | ↑6.00% ↓2.80% | 3.61% 2.08% |
| Initiation & Engagement of Alcohol & Other Drug Dependance Treatment – Engagement | 16.33% | ↑24.09% | 14.37% |
| Ages 13-17 | FO 070/ | L F2 O20/ | FC 130/ |
| Follow-up After Hospitalization Mental Illness - 7 Day Follow-up (Ages 6 to 17) | 58.87% | ↓ 53.03% | 56.13% |
| Follow-up after Mental Illness-7 Day Follow-up (ages 18-64) Total FU for Children Prescribed ADHD Med - Initiation Phase | 42.85% | ↓ 34.65% | 46.02% |
| | 44.78% | ↓ 44.30% | 44.17% |
| FU for Children Prescribed ADHD Med - Continuation and Maintenance Phase | 56.70% | ↓ 53.39% | 57.53% |
| Follow-up After ED visit for Mental Illness - 7 day | 76.33% | ↓ 70.69% | 64.39% |
| Follow-up After ED visit for Mental Illness - 30 day | 84.64% | ↓ 82.66% | 76.87% |
| First Line Psychosocial Care for Children/Adolescents on Antipsychotics Total | 79.15% | ↓ 76.16% | 69.30% |
| Comprehensive Diabetes Care | 26.000/ | A 27 060/ | 25 520/ |
| HbA1c Poor Control (lower is better) ->9.0% | 36.98% | ↑ 37.96% | 35.52% |
| HbA1c Control for Patients with Diabetes - <8.0% | 53.28% | <u>↑54.74%</u> | 54.26% |
| Retinal Eye Exam Performed | 56.93% | ↓ 51.34% | 56.51% |
| Blood Pressure Control (<140/90 mm Hg) | 63.75% | ↓ 63.50% | 67.40% |
| Kidney Health Evaluation for Patients with DM | 30.97% | ↑31.44% | NA 50.530/ |
| Received Statin Therapy for Patients with DM | 66.65% | ↓ 64.36% | 69.52% |
| Cardiovascular Conditions | | A | |
| Controlling High Blood Pressure | 61.31% | ↑ 62.29% | 65.10% |
| Statin Therapy for Patients with CV Disease | 81.52% | ↓ 80.45% | 76.22% |
| Respiratory Conditions | | | |
| Annual Monitoring for Patients on Persistent Medications - Total | 81.32% | 1 87.90% | 87.18% |
| Pharmacology Management of COPD Exacerbation: Corticosteroid | 74.23% | ↑ 73.63% | 76.87% |
| Pharmacology Management of COPD Exacerbation: Bronchodilator | 86.25% | ↓ 85.03% | 88.67% |
| Well Child Visits within 15 months of life – 6 or more | 51.81% | ↑ 60.47% | 61.19% |
| Well Child Visits Age 15-30 months of Life | 57.24% | ↑ 59.74% | 72.24% |
| Child & Adolescent Well-Care Visits, 3-11 years | 49.16% | ↑ 50.41% | 64.19% |
| Child & Adolescent Well-Care Visits, 12-17 years | 43.16% | 1 43.79% | 58.47% |

BUCKEYE HEALTH PLAN

MEDICAID ANNUAL HEDIS RATES 2022-2023

| Medicaid HEDIS® Measures, continued | HEDIS 2022 (MY2021) | HEDIS 2023 (MY2022) | Goal |
|---|---------------------------|---------------------------|--------|
| Child & Adolescent Well-Care Visits, 18-21 years | 21.63% | ↑ 21.72% | 31.88% |
| Child & Adolescent Well-Care Visits, Total | 42.84% | 1 43.38% | 53.83% |
| BMI Percentile for Children/Adolescents | 73.72% | ↓ 69.34% | 84.43% |
| Lead Screening in Children | 57.16% | ↑ 60.07% | 72.67% |
| Annual Dental Visit | 34.77% | ↑ 36.56% | 55.89% |
| Childhood Immunizations - Combo 3 | 59.37% | ↑ 62.53% | 68.86% |
| Childhood Immunizations - Combo 10 | 61.63% | ↓ 25.06% | 42.09% |
| Immunizations for Adolescents - Combo 1 | 73.48% | ↑ 74.45% | 85.40% |
| Immunizations for Adolescents - HPV | 29.44% | ↓ 27.98% | 42.58% |
| Immunizations for Adolescents - Combo 2 | 27.98% | ↓ 26.76% | 41.12% |
| Timeliness of Prenatal Care | 84.91% | ↓ 82.73% | 88.26% |
| Postpartum Care | 76.89% | ↓ 71.29% | 81.27% |
| Breast Cancer Screening | 53.75% | ↓ 52.51% | 56.52% |
| Cervical Cancer Screening, Total | 55.96% | ↑ 58.15% | 62.53% |
| Chlamydia Screening in Women, Total | 55.28% | ↓ 54.71% | 62.65% |
| Adult Access to Preventive Care - Total | 77.07% | ↓ 75.25% | 80.86% |
| Ambulatory Care - Emergency Department Visits (lower is better) | 67.33% | ↑ 77.87% | 51.60% |

[↑]Increase/↓Decrease over prior year